ClearBand SAFE DNA Gel Stain Solution, 20,000x

 $1 \, \mathrm{ml}$

Cat No: SDGS1

Shipping : Ship at ambient temperature.

Storage : Stable for at least 1 year when stored at 4°C.

General Information

ClearBand SAFE DNA Gel Stain Solution, 20,000x, is an aqueous fluorescent nucleic acid gel stain as a safer alternative to traditional ethidium bromide (EtBr) to be used at 1x concentration in agarose or polyacrylamide gels.

One vial (1 ml) of *ClearBand* SAFE DNA Gel Stain Solution, 20,000x, can be used to prepare at least 200 precast mini gels. *ClearBand* SAFE DNA Gel Stain Solution, 20,000x is compatible with downstream applications including gel extraction and cloning.

Key Features;

Safety: No toxicity, no mutagenicity and no carcinogenicity.

Eco-friendly: Safe to dispose in the drain. No hazard wastes.

Affordable: Low cost. An affordable alternative to competing dyes.

Compatibility: Compatible with instruments that are suitable for detection of gels with EtBr. Suitable for blue LED detection.

Protocol

The staining protocol for *ClearBand* SAFE DNA Gel Stain Solution, 20,000x is similar to that for EtBr.

Note: Briefly vortex *ClearBand* SAFE DNA Gel Stain Solution, 20,000x before use.

- 1. Prepare agarose gel solution in a flask using your standard protocol. Heat the agarose solution until the solution is completely clear.
- 2. Add 5µl of *ClearBand* SAFE DNA Gel Stain Solution, 20,000x per 100 ml of the gel before casting the gel. Swirl the flask gently to mix the solution and avoid producing air bubbles. *ClearBand* SAFE DNA Gel Stain Solution, 20,000x can be added while the gel solution is still hot.
- **3.** Pour the solution from step 2 into the gel tray, insert the combs and allow the agarose gel to cool until it solidifies.

Note: The thickness of gel is recommended to be less than 0.5 cm to increase the sensitivity.

- **4.** Load samples on the gel and perform electrophoresis.
- **5.** Detect the bands under UV illumination or blue LED.